


BRL  
1149  
c.1A

REFERENCE COPY

GA



REPORT NO. 1149  
SEPTEMBER 1961

AN INTEGRAL EQUATION OCCURRING IN  
A PROBLEM OF SUBSIDENCE

J. H. GIESE

TECHNICAL LIBRARY  
U. S. ARMY ORDNANCE  
ABERDEEN PROVING GROUND, MD.  
ORDEG-TL

Department of the Army Project No. 503-06-002  
Ordnance Management Structure Code No. 5010.11.812  
**BALLISTIC RESEARCH LABORATORIES**



**ABERDEEN PROVING GROUND, MARYLAND**

BRL  
1149

u

C. 1A

OTICE

An integral equation occurring

this report from ASTIA.

Quali: in a problem of subsidence

JUN 22 1976

*Rosalia Zerst (sp)*

RECEIVED  
U. S. AIR FORCE  
ARWEDSEN ENGINEERING  
CAMP 11

Library Bureau

BALLISTIC RESEARCH LABORATORIES

REPORT NO. 1149

SEPTEMBER 1961

AN INTEGRAL EQUATION OCCURRING IN A PROBLEM OF SUBSIDENCE

J. H. Giese

TECHNICAL LIBRARY  
U S. ARMY ORDNANCE  
ABERDEEN PROVING GROUND, MD.  
ORDBG-TL

Department of the Army Project No. 503-06-002  
Ordnance Management Structure Code No. 5010.11.812

ABERDEEN PROVING GROUND, MARYLAND

BALLISTIC RESEARCH LABORATORIES

REPORT NO. 1149

JHGiese/bjw  
Aberdeen Proving Ground, Md.  
September 1961

AN INTEGRAL EQUATION OCCURRING IN A PROBLEM OF SUBSIDENCE

ABSTRACT

The linear boundary value problem that was solved by numerical methods in BRL Report No. 1131 has been reformulated as a linear integral equation and solved by Laplace transform methods.

Page intentionally blank

Page intentionally blank

Page intentionally blank

## INTRODUCTION

As a result of mining operations cavities are produced into which the overburden may eventually collapse. Following collapse there will be a period during which the fallen material subsides. It would be extremely difficult to treat this motion in a way that takes into account all the geometrical and physical peculiarities of a particular situation. However, Leser and Jenike <sup>(1)</sup> have proposed a semi-empirical idealized one-dimensional model of subsidence which is based on the following system of linear partial differential equations.

$$\partial u / \partial T = -\alpha \partial y / \partial T \quad (1.1)$$

$$\partial y / \partial t = u \quad (1.2)$$

Here  $y(t, T)$  is the height above the top surface of the cavity at time  $t$  of the layer of material through which the plane of disturbance passed at time  $T$ ;  $u(t, T)$  is the velocity of this layer; and

$$\alpha(t, T) = K \left[ 1 + K(t - T) \right]^{-1} - \left[ 1 + t - T \right]^{-1} \quad (1.3)$$

is an empirical function, in which the constant  $K > 1$ . By eliminating  $y(t, T)$  between (1.1) one obtains for  $u(t, T)$  the linear hyperbolic partial differential equation

$$\frac{\partial^2 u}{\partial t \partial T} = \frac{-2K}{1 + K(t - T)} \frac{\partial u}{\partial T} \quad (1.4)$$

Boundary conditions for (1.4) were determined by the following considerations. Let  $\gamma_s$  be the original density of the undisturbed overburden, and  $\gamma_o$   $\gamma_s$  the density immediately after passage of the disturbance, both being assumed constant. Then conservation of matter at the wave front  $y(t, t)$  implies

$$\gamma_s \, dy(T, T)/dT = \gamma_o \left[ dy(T, T)/dT - u(T, T) \right] \quad (1.5)$$

By (1.1) - (1.3) this can be rewritten as

$$\partial u(T, T)/\partial T = \gamma_s (K - 1) (\gamma_s - \gamma_o)^{-1} u(T, T) \quad (1.6)$$

On  $T = 0$  it was assumed on empirical grounds that for some constant  $\epsilon > 0$

$$u(t, 0) = -\epsilon \gamma_s (K - 1) (\gamma_s - \gamma_0)^{-1} (1 + Kt)^{-2} \quad (1.7)$$

Leser and Jenike solved this problem originally by numerical integration. However, as we shall show in the following sections:

(i) An integral representation (2.6) can be found for  $u(t, T)$  in terms of  $u(t, t)$ ;

(ii)  $u(t, t)$  satisfies a Volterra integral equation (2.8) of the second kind, of convolution type, that depends on the single parameter

$$a = (1 - K^{-1}) (1 - \gamma_0 \gamma_s^{-1})^{-1} > 0 \quad (1.8)$$

(iii)  $\lim_{t \rightarrow \infty} u(t, t) = 0$  or  $\infty$  as  $a \leq 1$  or  $a > 1$ .

(iv)  $1 \leq (1 + t)^2 u(t, t) \leq m^{-2}(a)$  for some  $m(a)$ ,  $a < 1$ .

(v)  $u(t, t)$  is a strictly increasing function of  $a$  for fixed  $t$ .

(vi)  $u(t, t)$  is a strictly decreasing function of  $t$  for fixed  $a$  such that  $0 < a \leq 1$ .

(vii) The wave-front  $y(t, t)$  advances only a finite distance if  $a < 1$ .

The author wishes to acknowledge his indebtedness to Dr. T. Leser for calling his attention to this problem, and to Dr. W. C. Taylor for suggestions concerning its treatment by Laplace transform theory.

#### REFORMULATION-REDUCTION TO AN INTEGRAL EQUATION

If we let

$$w(t, T) = -\frac{(\gamma_s - \gamma_0)}{\epsilon \gamma_s (K - 1)} u\left(\frac{t}{K}, \frac{T}{K}\right) \quad (2.1)$$

then (1.5) - (1.7) become

$$\frac{\partial^2 w}{\partial t \partial T} = -\frac{2}{1 + t - T} \frac{\partial w}{\partial T} \quad (2.2)$$

$$\frac{\partial w(T, T)}{\partial T} = a w(T, T) \quad (2.3)$$

$$w(t, 0) = (1 + t)^{-2} \quad (2.4)$$

where

$$a = \gamma_s (K - 1)/K (\gamma_s - \gamma_0) > 0 \quad (2.5)$$

Now (2.2) to (2.4) imply

$$w(t, T) = \frac{1}{(1 + t)^2} + a \int_0^T \frac{w(r, r) dr}{(1 + t - r)^2} \quad (2.6)$$

and if we let

$$v(t) = w(t, t) \quad (2.7)$$

then by (2.6)

$$v(t) = \frac{1}{(1 + t)^2} + a \int_0^t \frac{v(r) dr}{(1 + t - r)^2} \quad (2.8)$$

Thus  $v(t)$  is defined as a solution of a Volterra integral equation of the second kind of convolution type. The standard theorem for such equations <sup>(2)</sup> asserts the existence of  $v(t)$ , and in fact yields

$$(1 + t)^{-2} \leq v(t) \leq e^{at} (1 + t)^{-2} \quad (2.9)$$

Also by (2.6)

$$(1 + t)^{-2} \leq w(t, T) \leq w(t, t) = v(t) \quad (2.10)$$

and, actually,  $w(t, T)$  is a strictly increasing function of  $T$  for fixed  $t = T$ . Since the positive function  $w(t, T)$  is dominated by  $v(t)$ , in order to provide a basis for estimating  $w$  it will suffice to consider only  $v(t)$ .

#### APPLICATION OF LAPLACE TRANSFORMS

By (2.9)  $v(t)$  is of exponential type. Accordingly its Laplace transform



$$V(s) = \int_0^{\infty} e^{-st} v(t) dt \quad (3.1)$$

is a regular function of  $s$  for  $\operatorname{Re} s > a$ . Also

$$f(t) = (1+t)^{-2} \quad (3.2)$$

is bounded, and its Laplace transform

$$F(s) = \int_0^{\infty} e^{-st} f(t) dt \quad (3.3)$$

is regular for  $\operatorname{Re} s > 0$ . Since

$$(e^{-s} F(s))' = e^{-s}/s \quad (3.4)$$

and since

$$e^{-s} F = (e^{-s} F)' = 0 \text{ for } \operatorname{Re} s = \infty,$$

then (3.4) implies, after an integration by parts, that

$$F(s) = 1 - se^s \int_s^{\infty} r^{-1} e^{-r} dr \quad (3.5)$$

which is regular in the entire  $s$ -plane except for a slit along the negative real-axis.

In the region where both  $V(s)$  and  $F(s)$  exist the Laplace transform of (2.8)

$$V(s) = F(s) + aF(s) V(s)$$

and then (3.5) and

$$V(s) = F(s) / [1 - a F(s)] \quad (3.6)$$

define the analytic continuation of  $V(s)$  into the slit  $s$ -plane. To determine the nature and location of the singularities of  $V$ , first

consider  $F$ . If we move the point  $s$  around a closed path that contains  $s = 0$  we obtain from (3.5)

$$F(se^{2\pi i}) = F(s) + 2\pi i s^s$$

Accordingly

$$F(s) = H(s) + se^s \log s \quad (3.7)$$

where  $H(s)$  is an entire function. Thus  $F(s)$  has a logarithmic branch point at  $s = 0$ . Since  $F$  is real on the positive real axis,  $H(s)$  is real on the entire real axis. Now

$$\operatorname{Im} F(s) = \begin{cases} 0 & \text{for } s \geq 0 \\ \pm se^s & \text{for } s \leq 0 \end{cases}$$

where the upper (lower) sign refers to the upper (lower) side of the slit. By the maximum principle for harmonic functions  $\operatorname{Im} F(s)$  is negative (positive) on the upper (lower) half of the  $s$ -plane. Consequently the roots of

$$1 - aF(s_0) = 0 \quad (3.8)$$

can lie only on the positive real axis or at  $s = 0$ . Since  $F(0) = 1$ , and since for  $s \geq 0$

$$F'(s) = - \int_0^\infty e^{-st} t f(t) dt < 0$$

(3.8) has a single simple root  $s_0 > 0$  if and only if  $a > 1$ ; and  $s_0 = 0$  if and only if  $a = 1$ . Thus, if  $a > 1$ , then  $V(s)$  has a single simple pole at  $s_0$  in addition to a branch point at 0; if  $0 < a < 1$  it has only a branch point.

The complex inversion formula yields

$$v(t) = (2\pi i)^{-1} \int_{x-1-i\infty}^{x+1+i\infty} e^{ts} v(s) ds \quad (3.9)$$

where  $x > s_0$ . An estimate that can be used to bound  $|v(s)|$  can be found by integrating (4.5) twice by parts to obtain

$$F(s) = s^{-1} - 2s^{-2} + 6se^s \int_s^\infty r^{-4} e^{-r} dr$$

If the path of integration of  $F(s)$  is taken to be the half-line

$$|s| \leq r \leq \infty \text{ and the circular arc } r = |s| e^{i\theta}, 0 \leq \theta \leq \phi \leq \pi,$$

then

$$|F(s) - s^{-1} + 2s^{-2}| \leq 6 |s|^{-3} e^{-2|s| \sin^2 \phi/2} + 6 |s|^{-2} \quad (3.10)$$

By standard techniques and application of (3.10) the path of integration in (3.9) can be deformed into a circle  $C$  about  $s_0$  (if  $a > 1$ ) and a path that traverses both sides of the negative real axis. The circle contributes

$$(2\pi i)^{-1} \int_C \frac{e^{st} F(s) ds}{1 - a F(s)} = - e^{s_0 t} F(s_0) / a F'(s_0), a > 1 \quad (3.11)$$

where, by (3.5) and (3.9)

$$F(s_0) = 1/a, \quad a F'(s_0) = (1 + s_0^{-1})(1 - a) + a \quad (3.12)$$

The negative real axis contributes

$$I(t) = (2\pi i)^{-1} \int_{-\infty}^0 e^{tx} (V^- - V^+) dx$$

where  $V^-$  ( $V^+$ ) designates values on the lower (upper) side of the axis

and  $x = \operatorname{Re} s = s$ . Now write

$$V^- - V^+ = F^- / (1 - aF^-) - F^+ / (1 - aF^+) = (F^- - F^+) / (1 - aF^-)(1 - aF^+)$$

By (3.7)  $F^+ = \overline{F^-}$  and  $F^- - F^+ = -2\pi i x e^x$ , whence

$$I(t) = - \int_{-\infty}^0 x e^{(1+t)x} |1 - aF^-(x)|^{-2} dx > 0. \quad (3.13)$$

If  $a \neq 1$ , then  $|1 - aF^-| \neq 0$  on  $-\infty \leq x \leq 0$ , since  $F^-(-\infty) = 0$

and  $F^-(0) = 1$ . Then if we let

$$m(a) = \min |1 - aF^-| > 0$$

we find

$$0 < I(t) < -m^{-2} \int_{-\infty}^0 x e^{(1+t)x} dx = m^{-2}(1+t)^{-2}, \quad a \neq 1. \quad (3.14)$$

Now we can prove the assertions of Section 1. If  $a > 1$ , then

$$v(t) = I(t) - e^{s_0 t} F(s_0) / aF'(s_0)$$

which implies part of (iii). If  $a < 1$ , then  $v(t) = I(t)$ , and (3.14) implies another part of (iii). In conjunction with  $v = (1+t)^{-2}$  (3.14) yields (iv).

The case  $a = 1$  remains to be treated. First note that by (3.7)  $F(0) = 1$  implies  $H(0) = 1$ . Now let

$$H(s) = 1 + \sum_{n=1}^{\infty} a_n s^n$$

where the coefficients  $a_n$  are real. Then

$$1 - F^- = - \sum_{n=1}^{\infty} a_n x^n - x e^x [\log(-x) - \pi i]$$

and

$$|1 - F^-|^2 = x^2 \left\{ \left[ \sum_{n=1}^{\infty} a_n x^{n-1} + e^x \log(-x) \right]^2 + \pi^2 e^{2x} \right\} \quad (3.15)$$

The convergence of  $I$  at its upper limit follows from

$$- \int_{-b}^0 \frac{e^{(t-1)x} dx}{x \log^2(-x)} \leq -N \int_{\log b}^{-\infty} u^{-2} du = -N/\log b \quad (3.16)$$

where  $0 < b < 1$ , and  $N = \max(1, e^{(1-t)b})$ . The convergence of  $I$  at its lower limit follows from  $|1 - F^-|^2 = O(x^2 e^{2x})$  for  $x \rightarrow -\infty$ . Now we can easily deduce

$$\lim_{t \rightarrow \infty} v(t) = 0 \text{ if } a = 1.$$

Proof: Write

$$I = \int_{-\infty}^{-b} + \int_{-b}^0 x e^{(1+t)x} |1 - F^-|^{-2} dx = I_1 + I_2$$

and suppose  $t > 1$ . First, choose  $b < 1$  so small that

$$|1 - F^-|^2 \geq 0.5 x^2 \log^2(-x) \text{ in } I_2. \text{ Then choose } \theta > 0 \text{ and reduce } b,$$

if necessary, so that  $-2/\log b < \theta$ . Then by (3.16)  $0 < I_2 < \theta$  for all

$t > 1$ . Next, if we let

$$\int_{-\infty}^{-b} x |1 - F^-|^{-2} dx = P$$

then

$$I_1 = e^{-(1+t)b} P,$$

and for

$$t > 1 - b^{-1} \log (\Theta/P)$$

we have  $0 < I_1 < \Theta$ . Finally, for such  $b$  and  $t$ ,  $0 < I < 2\Theta$ .

The standard iterative technique for solving (2.8) would yield a power series in  $a$  with coefficients that are positive functions of  $t$ . Thus  $v(t)$  is a monotone function of  $a$  for all  $t > 0$ . This and the representation  $v(t) = I(t)$  for  $0 < a \leq 1$ , and (3.13) imply (v) and (vi).

Finally, let us consider the motion of the wave-front  $y = y(t, t)$ . By (1.5), (2.1), and (2.7)

$$dy(t, t)/dt = \epsilon(K-1) \gamma_0 \gamma_s (\gamma_s - \gamma_0)^{-2} v(Kt)$$

and if  $y(0, 0) = 0$ , then

$$y(t, t) = \epsilon K^{-1}(K-1) \gamma_0 \gamma_s (\gamma_s - \gamma_0)^{-2} \int_0^{Kt} v(s) ds \quad (3.17)$$

By (iv), if  $a \neq 1$ .

$$1 \leq (1+t^{-1}) \int_0^t v(s) ds = m^{-2}(a),$$

which implies (vii).

*J. H. Giese*

J. H. GIESE

#### REFERENCES

1. Leser, T., and Jenike, A. Numerical Solutions of the Underground Subsidence Problem. BRL Report No. 1131 (1961); to be submitted to Mining Engineering Monthly.
2. Tricomi, F. G. Integral Equations. New York, 1957.

# DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>	<u>No. of Copies</u>	<u>Organization</u>
10	Commander Armed Services Technical Information Agency ATTN: TIPCR Arlington Hall Station Arlington 12, Virginia	1	Commander U.S. Naval Weapons Laboratory ATTN: Dr. Cohen Dahlgren, Virginia
1	Commanding Officer Diamond Ordnance Fuze Laboratories ATTN: Technical Information Office, Branch 041 Washington 25, D.C.	1	Commander Air Proving Ground Center ATTN: PGAPI Eglin Air Force Base, Florida
2	Commanding General ATTN: Technical Library White Sands Missile Range, New Mexico	1	U.S. Department of the Interior U.S. Bureau of Mines 4800 Forbes Street Pittsburgh 13, Pennsylvania
1	Commanding Officer Army Research Office (Durham) Box CM, Duke Station Durham, North Carolina	1	Hercules Experimentation Station Wilmington, Delaware
1	Army Research Office Arlington Hall Station Arlington, Virginia	1	Vitro Laboratories ATTN: Library 200 Pleasant Valley Way West Orange, New Jersey
1	Chief of Ordnance ATTN: ORDTB - Bal Sec Department of the Army Washington 25, D.C.	1	Cornell University Department of Mining Engineering Ithaca, New York
3	Chief, Bureau of Naval Weapons ATTN: DIS-33 Department of the Navy Washington 25, D.C.	1	Ohio State University Department of Mining Engineering Columbus 8, Ohio
2	Commander Naval Ordnance Laboratory ATTN: Library White Oak, Silver Spring 19, Maryland	1	Princeton University Department of Mining Engineering ATTN: Librarian Princeton, New Jersey
2	Commander U.S. Naval Ordnance Test Station ATTN: Technical Library China Lake, California	1	Purdue University Department of Mining Engineering Lafayette, Indiana
		1	Southwest Research Institute ATTN: F.A. Warren 8500 Culebra Road San Antonio 6, Texas



# DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>	<u>No. of Copies</u>	<u>Organization</u>
1	University of Kentucky Department of Mathematics ATTN: Dr. A.W. Goodman Lexington, Kentucky	10	Commander British Army Staff British Defence Staff (W) ATTN: Reports Officer 3100 Massachusetts Avenue, N.W. Washington 8, D.C.
1	Dr. A.S. Galbraith Army Research Office Box CM, Duke Station Durham, North Carolina		Of Interest to:
1	Dr. I.R. Hershner Army Research Office Arlington Hall Station Arlington, Virginia		Dr. H.R. Fehling 7 Albion Gate London, England
1	Dr. W.H. Pell National Bureau of Standards 232 Dynamometer Building Washington 25, D.C.		Professor C.J. Tranter Military College of Science Shrivenham, England
1	Dr. R. Langer U.S. Army Mathematics Center Madison, Wisconsin	4	Defence Research Member Canadian Joint Staff 2450 Massachusetts Avenue, N.W. Washington 8, D.C.

AD Accession No.  
Ballistic Research Laboratories, APG  
AN INTEGRAL EQUATION OCCURRING IN A  
PROBLEM OF SUBSIDENCE  
J. H. Giese

UNCLASSIFIED

BRL Report No. 1149 September 1961

DA Proj No. 503-06-002, OMSC No. 5010.11.812  
UNCLASSIFIED Report

The linear boundary value problem that was solved by numerical methods in PRL Report No. 1151 has been reformulated as a linear integral equation and solved by Laplace transform methods.

AD Accession No.  
Ballistic Research Laboratories, APG  
AN INTEGRAL EQUATION OCCURRING IN A  
PROBLEM OF SUBSISTENCE  
J. H. Giese

UNCLASSIFIED

Mathematics - Application  
Subsidence Problem -  
Numerical Solution

BRL Report No. 1149 September 1961

DA Proj No. 503-06-002, OMSC No. 5010.11.812  
UNCLASSIFIED Report

The linear boundary value problem that was solved by numerical methods in ERL Report No. 1131 has been reformulated as a linear integral equation and solved by Laplace transform methods.

AD Accession No.  
Ballistic Research Laboratories, APG  
AN INTEGRAL EQUATION OCCURRING IN A  
PROBLEM OF SUBSIDENCE  
J. H. Giese

UNCLASSIFIED  
Issues - Applications  
the Problem -  
Real Solution

BRL Report No. 1149 September 1961

DA Proj No. 503-06-002, OMSC No. 5010.11.812  
UNCLASSIFIED Report

The linear boundary value problem that was solved by numerical methods in BRL Report No. 1151 has been reformulated as a linear integral equation and solved by Laplace transform methods.

AD Accession No.  
Ballistic Research Laboratories, APG  
AN INTEGRAL EQUATION OCCURRING IN A  
PROBLEM OF SUBSIDENCE  
J. H. Giese

UNCLASSIFIED

Mathematics - Application  
Subsidence Problem -  
Numerical Solution

BRL Report No. 1149 September 1961

DA Proj No. 503-06-002, OMSC No. 5010.11.812  
UNCLASSIFIED Report

The linear boundary value problem that was solved by numerical methods in BRL Report No. 1131 has been reformulated as a linear integral equation and solved by Laplace transform methods.